



locessos leziol

NUKNIT-BUPIVACAINE						
		OGEN UN	NDER NITR ITS USE	STORE UNDER NITROGEN UNTIL	01	
		(0	DRY SAMPLES	DRY		
				COOL UNDER NITROGEN FOR OVERNIGHT	OL UNDER NITR FOR OVERNIGHT	202
	띨	DRIED NUKNIT-BUPIVACAINE	NUKNIT-	DRIED		
			Or .	DRY AT 50 C FOR AN HOUR	DRY AT AN	
		CAINE	NUKNIT-BUPIVACAINE	NOKN		
				ADD 10% BUPIVACAINE SOLUTION) 10% BU SOLUTION	ADE
	NUKNIT PIECES (1.0" X 0.6") ~80 mg	" X 0.6")	SES (1.0	JKNIT PIE(ž	



--- NKM F1 UNCOATED
---- NKM F2 COATED (LOW LEVEL)
--- NKM F3 COATED (HIGH LEVEL)

F.I.G. 6



TODESSOS JESTOI

NuKnit + 2 LAYERS OF FILM (OR)

NuKnit + 2 LAYERS OF FILM + 2LAYERS OF NuKnit

NuKnit + 2 LAYERS OF FILM (OR) NuKnit + 2 LAYERS OF FILM + 2LAYERS OF NuKnit

FIG. 8